

Sports injuries and illnesses during the London Summer Olympic Games 2012

Lars Engebretsen,^{1,2,3} Torbjørn Soligard,¹ Kathrin Steffen,³ Juan Manuel Alonso,⁴ Mark Aubry,⁵ Richard Budgett,¹ Jiri Dvorak,^{6,7} Manikavasagam Jegathesan,^{8,9,10} Willem H Meeuwisse,¹¹ Margo Mountjoy,¹² Debbie Palmer-Green,¹³ Ivor Vanhegan,¹⁴ Per A Renström¹⁵

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/bjsports-2013-092380>).

For numbered affiliations see end of article.

Correspondence to

Torbjørn Soligard, Medical & Scientific Department, International Olympic Committee, Lausanne 1007, Switzerland; torbjorn.soligard@olympic.org

Accepted 27 February 2013

ABSTRACT

Background The Olympic Movement Medical Code encourages all stakeholders to ensure that sport is practised without danger to the health of the athletes. Systematic surveillance of injuries and illnesses is the foundation for developing preventive measures in sport. **Aim** To analyse the injuries and illnesses that occurred during the Games of the XXX Olympiad, held in London in 2012.

Methods We recorded the daily occurrence (or non-occurrence) of injuries and illnesses (1) through the reporting of all National Olympic Committee (NOC) medical teams and (2) in the polyclinic and medical venues by the London Organising Committee of the Olympic and Paralympic Games' (LOCOG) medical staff.

Results In total, 10 568 athletes (4676 women and 5892 men) from 204 NOCs participated in the study. NOC and LOCOG medical staff reported 1361 injuries and 758 illnesses, equalling incidences of 128.8 injuries and 71.7 illnesses per 1000 athletes. Altogether, 11% and 7% of the athletes incurred at least one injury or illness, respectively. The risk of an athlete being injured was the highest in taekwondo, football, BMX, handball, mountain bike, athletics, weightlifting, hockey and badminton, and the lowest in archery, canoe slalom and sprint, track cycling, rowing, shooting and equestrian. 35% of the injuries were expected to prevent the athlete from participating during competition or training. Women suffered 60% more illnesses than men (86.0 vs 53.3 illnesses per 1000 athletes). The rate of illness was the highest in athletics, beach volleyball, football, sailing, synchronised swimming and taekwondo. A total of 310 illnesses (41%) affected the respiratory system and the most common cause of illness was infection (n=347, 46%).

Conclusions At least 11% of the athletes incurred an injury during the games and 7% of the athletes' an illness. The incidence of injuries and illnesses varied substantially among sports. Future initiatives should include the development of preventive measures tailored for each specific sport and the continued focus among sport bodies to institute and further develop scientific injury and illness surveillance systems.

INTRODUCTION

According to the Olympic Charter, the codification governing the organisation and operation of the Olympic Movement, one of the main roles of the International Olympic Committee (IOC) is to encourage and support measures protecting the health of athletes.¹ Systematic monitoring of injury

and illness trends over long periods of time provides epidemiological data that are indispensable to identify and subsequently reduce injuries and illnesses in high-risk sports and disciplines.² Thus, to maximise the health protection and benefits of elite athletes, as well as to minimise the direct and indirect costs associated with injury and illness, the early identification of athletes at a high risk and subsequent development and introduction of effective prevention tools are significant goals for the IOC.

As early as 1998, the Fédération Internationale de Football Association (FIFA) started to survey all injuries incurring during their competitions,^{3–8} and other major sports federations followed their lead.^{9–16} Based on these experiences, a group of experts, convened by the IOC, developed an injury surveillance system for multi-sport events.¹⁷ It was successfully implemented for the first time in the 2008 Beijing (surveillance of injuries only)¹⁸ and in the 2010 Vancouver Olympics (surveillance of both injuries and illnesses).¹⁹ We further developed and optimised it for the Games of the XXX Olympiad, held in London in 2012 (hereafter referred to as the 'London Games').

With almost 11 000 athletes from 204 countries, the London Games was one of the largest sports events ever. The aim of the present paper is to analyse the injuries and illnesses that occurred in London, with the long-term aim to enable the National Olympic Committees (NOCs) and International Federations (IF) to improve their work on protection of their athletes' health. Practical implications and suggestions for further research to protect the athletes' health are provided.

METHODS

We employed the IOC injury and illness surveillance system for multisport events to record injuries and illnesses in the present study.^{18, 19} First, we asked all NOC head physicians to report the daily occurrence (or non-occurrence) of newly sustained injuries and illnesses on a standardised form. Second, we retrieved the same information on all athletes treated for injuries and illnesses in the polyclinic and medical venues by the London Organising Committee of the Olympic and Paralympic Games' (LOCOG) medical staff.

Implementation

We informed the NOCs about the study in a letter sent 4 months in advance of the Olympic Games.

To cite: Engebretsen L, Soligard T, Steffen K, *et al.* *Br J Sports Med* Published Online First: [please include Day Month Year] doi:10.1136/bjsports-2013-092380

The day before the opening of the Games, we invited the medical staff of the NOCs to a meeting covering the details of the study. In this meeting, we also started the distribution of the daily injury and illness report forms, as well as an instructional booklet detailing the study protocol.

Throughout the data collection, we recorded the response rate of NOCs having more than 30 participating athletes, and frequently visited these to encourage continuous medical reporting throughout the games.

Definition of injury and illness

We defined injuries and illnesses as new (pre-existing, not fully rehabilitated conditions were not recorded) or recurring (athletes having returned to full participation after a previous condition) musculoskeletal complaints or concussions (injuries) or illnesses incurred during competition or training during the London Olympic Games (27 July–12 August 2012) receiving medical attention, regardless of the consequences with respect to absence from competition or training.¹⁷ Injuries and illnesses occurring in football were recorded 25 July onwards, the opening date of the football competition. In cases where a single incident caused multiple injury types or affected multiple body parts, we recorded only the most severe diagnosis.^{17 19 20}

Injury and illness report form

The injury and illness report form was identical to the one introduced in the XXI Olympic Winter Games in Vancouver in 2010.¹⁹ With respect to injuries, we recorded the following information: accreditation number, sport and event, whether the injury occurred during competition or training, date and time, body part, type, cause and estimated time lost from competition or training. Likewise, we recorded the following information for illnesses: accreditation number, sport and event, date, diagnosis, affected system, main symptom(s), cause and estimated time loss.

We provided instructions on how to complete the form correctly in the instructional booklet. We distributed the injury and illness report forms to all NOCs in the following languages: English, French, Arabic, Chinese, German, Japanese, Russian and Spanish.

Confidentiality and ethical approval

We recorded and utilised the athlete's accreditation number to control for duplicates resulting from certain athletes being treated for the same condition by both the NOC and the LOCOG medical teams. In cases of duplicates, we retained the NOC data. We also used the accreditation number to query the IOC athlete database for the athletes' age, gender, nationality and sport. We treated all information with strict confidence, and anonymised our medical database at the end of the Games.

The study was approved by the medical research ethics committee of the South-Eastern Norway Regional Health Authority, Norway, and by the University College London Research Ethics Committee, UK.

Data analysis

We calculated the summary measure of injury and illness incidence (i) according to the formula $i=n/e$, where n is the number of injuries or illnesses during competition, training or in total during the study period and e the respective number of exposed athletes. We calculated CIs of the relative risk (RR) of the number of injuries or illnesses between two groups by a simple Poisson model, assuming constant hazard per group. We present

injury and illness incidences as means and rate ratios with 95% CIs. We regarded two-tailed p values ≤ 0.05 as significant.

RESULTS

In total, 10 568 athletes took part in the London Olympic Games. Of these, 4676 were women (44%) and 5892 men (56%).

Incidence and distribution of injuries

Among these athletes, we recorded a total of 1361 injuries, equalling an overall injury rate of 128.8 injuries per 1000 registered athletes (table 1). On average, 11% of the athletes sustained at least one injury (n=1190). There were 114, 18 and 7 athletes with two, three and four injuries, respectively. The risk of an athlete to be injured was the highest in taekwondo, football, BMX, handball, mountain bike, hockey, weightlifting, athletics and badminton (15–39% of registered athletes were affected in each sport). The relative injury risk was the lowest for archery, canoe slalom and sprint, track cycling, rowing, shooting and equestrian (less than 5% of the athletes were injured). The injury rates in women (132.8 injuries per 1000 athletes (95% CI 122.4 to 143.3)) and men (121.0 (112.1 to 129.9)) was similar (RR=1.10 (0.97 to 1.22), p=0.11) (table 2). However, male athletes were at significantly higher risk of injury in taekwondo (RR=1.9 (1.1 to 3.5), p=0.03), whereas women were at higher risk of injury in football (RR=1.7 (1.2 to 2.2), p<0.001).

Severity, location and type of injuries

While two-thirds of the injuries were estimated to not result in any time loss from sport (n=879, 65%), a total of 482 (35%) injuries were expected to prevent the athlete from participating in competition or training. It was estimated that 246 (18%) injuries would result in an absence from sports for 1–3 days, 62 (5%) in an absence for 4–7 days, 105 (8%) in an absence for 8–28 days and 69 (5%) in an absence for more than 28 days.

A total of 174 injuries (13%) entailed an estimated absence from training or competition for more than 1 week (table 1). These injuries were 10 shoulder, elbow and knee dislocations (in hockey, football, judo, BMX and weightlifting); 38 muscle strains, of which 24 were thigh strains (mostly in athletics); 24 fractures (mostly in team sports; all body locations) and 6 stress fractures (4 in running events); 8 Achilles, knee and shoulder tendon ruptures (in athletics, badminton, handball and basketball); 47 ligament sprains (across all joints and sports) and 15 knee sprains, including 6 ACL and 1 PCL ruptures (in fencing, handball, judo, wrestling, badminton, table tennis, tennis and football). None of the six reported concussions (3 in football, 1 each in water polo, athletics and BMX) were classified as severe.

Mechanisms and circumstances of injury

The four most commonly reported injury mechanisms were overuse (n=346, 25%), non-contact trauma (n=275, 20%), contact with another athlete (n=197, 14%) and contact with a stationary object (n=164, 12%). Stratified by sport, overuse was the mechanism for 73% of all diving injuries, 34% of athletics and weightlifting injuries and 28% and 26% of swimming and fencing injuries, respectively. Non-contact trauma was a frequent mechanism in tennis (48% of all tennis injuries), beach volleyball and canoe sprint (43% each), fencing (39%), badminton (35%) and athletics (28%). In contrast, more than 40% of the injuries in basketball, football, handball and judo were caused by contact with another athlete. Of all overuse injuries, 234

Table 1 Rates of overall injuries, injuries leading to time loss (≥ 1 or >7 days of estimated absence), competition and training injuries and overall illnesses in the Olympic sports

Olympic sport	Athletes (n)	Injuries			Competition	Training	All illnesses
		All	≥ 1 day	>7 days			
Archery	128	2 (1.6)	0 (0.0)	0 (0.0)	0 (0.0)	1 (100.0)	10 (7.8)
Athletics	2079	368 (17.7)	145 (7.0)	59 (2.8)	133 (39.5)	204 (60.5)	219 (10.5)
Aquatics							
Diving	136	11 (8.1)	5 (3.7)	2 (1.5)	2 (25.0)	8 (75.0)	7 (5.1)
Swimming	931	50 (5.4)	7 (0.8)	1 (0.1)	13 (31.0)	29 (67.0)	68 (7.3)
Synchronised swimming	104	14 (13.5)	4 (3.8)	0 (0.0)	2 (20.0)	8 (80.0)	13 (12.5)
Water polo	260	34 (13.1)	13 (5.0)	0 (0.0)	26 (78.8)	7 (21.2)	21 (8.1)
Badminton	164	26 (15.9)	7 (4.3)	4 (2.4)	11 (47.8)	12 (52.3)	5 (3.0)
Basketball	287	32 (11.1)	10 (3.5)	7 (2.4)	21 (75.0)	7 (25.0)	9 (3.1)
Beach volleyball	96	12 (12.5)	2 (2.1)	0 (0.0)	6 (54.5)	5 (45.5)	18 (18.8)
Boxing	283	26 (9.2)	9 (3.2)	1 (0.4)	16 (72.7)	6 (27.3)	18 (6.4)
Canoe slalom	83	2 (2.4)	1 (1.2)	0 (0.0)	0 (0.0)	2 (100.0)	4 (4.8)
Canoe sprint	249	7 (2.8)	1 (0.4)	0 (0.0)	3 (50.0)	3 (50.0)	14 (5.6)
Cycling							
BMX	48	15 (31.3)	5 (10.4)	2 (4.2)	11 (73.3)	4 (26.7)	2 (4.2)
MTB	76	16 (21.1)	8 (10.5)	2 (2.6)	5 (31.3)	11 (68.7)	5 (6.6)
Road	210	19 (9.0)	7 (3.3)	2 (0.9)	14 (73.7)	5 (26.3)	7 (3.3)
Track	167	5 (3.0)	3 (1.8)	0 (0.0)	1 (20.0)	4 (80.0)	16 (9.6)
Equestrian	199	9 (4.5)	4 (2.0)	2 (1.0)	6 (100.0)	0 (0.0)	11 (5.5)
Fencing	246	23 (9.3)	10 (4.1)	2 (0.8)	10 (45.5)	12 (54.5)	13 (5.3)
Football	509	179 (35.2)	67 (13.2)	11 (2.2)	132 (74.2)	46 (25.8)	62 (12.2)
Gymnastics							
Artistic	195	15 (7.7)	8 (4.1)	4 (2.1)	8 (66.7)	4 (33.3)	5 (2.6)
Rhythmic	96	7 (7.3)	1 (1.0)	0 (0.0)	1 (16.7)	5 (83.3)	1 (1.0)
Trampoline	32	2 (6.3)	0 (0.0)	0 (0.0)	0 (0.0)	2 (100.0)	1 (3.1)
Handball	349	76 (21.8)	32 (9.2)	16 (4.6)	55 (75.3)	18 (24.7)	17 (4.9)
Hockey	388	66 (17.0)	25 (6.4)	10 (2.6)	44 (71.0)	18 (29.0)	29 (7.5)
Judo	383	47 (12.3)	22 (5.7)	12 (3.1)	26 (68.4)	12 (31.6)	16 (4.2)
Modern pentathlon	72	6 (8.3)	2 (2.8)	0 (0.0)	3 (60.0)	2 (40.0)	1 (1.4)
Rowing	549	18 (3.3)	2 (0.4)	0 (0.0)	7 (53.8)	6 (46.2)	40 (7.3)
Sailing	380	56 (14.7)	3 (0.8)	1 (0.3)	30 (62.5)	18 (37.5)	38 (10.0)
Shooting	390	15 (3.8)	4 (1.0)	0 (0.0)	1 (7.1)	13 (92.9)	17 (4.4)
Table tennis	174	11 (6.3)	7 (4.0)	2 (1.1)	7 (70.0)	3 (30.0)	12 (6.9)
Taekwondo	128	50 (39.1)	16 (12.5)	7 (5.5)	16 (33.3)	32 (66.7)	14 (10.9)
Tennis	184	21 (11.4)	7 (3.8)	4 (2.2)	14 (66.7)	7 (33.3)	4 (2.2)
Triathlon	110	16 (14.5)	8 (7.3)	3 (2.7)	11 (73.3)	4 (26.7)	7 (6.4)
Volleyball	288	20 (6.9)	7 (2.4)	3 (1.0)	11 (55.0)	9 (45.0)	8 (2.8)
Weightlifting	252	44 (17.5)	19 (7.5)	11 (4.4)	18 (45.0)	22 (55.0)	10 (4.0)
Wrestling	343	41 (12.0)	11 (3.2)	6 (1.7)	20 (62.5)	12 (37.5)	16 (4.7)
Total	10568	1361 (12.9)	482 (4.6)	174 (1.6)	684* (54.9)	561* (45.1)	758 (7.2)

Values are numbers (percentages) of injured or ill athletes, unless otherwise indicated.

*Information on training/competition is missing in 116 injuries.

(68%) were recorded with no estimated absence from competition or training.

The distribution of injuries during competition and training was similar, with 55% occurring in competition and 45% during training. Injuries during training and in competition differed significantly in all injury characteristics (location, type, mechanism and subsequent time loss from sport) and in terms of the different sports (table 1). Seventy-five per cent of all injuries in team sports occurred in competitions, whereas 60% of all injuries in athletics were sustained during training. Furthermore, of the more severe injuries, entailing an estimated absence from training or competition of more than 1 week, 81% occurred during competition.

Incidence and distribution of illnesses

Among the 10 568 athletes, a total of 758 illnesses were reported, resulting in an incidence of 71.7 illnesses per 1000 athletes (table 1). On average, 7% of the athletes incurred an illness. The incidence of illnesses was significantly higher in women compared to male athletes (86.0 vs 53.3 illnesses per 1000 athletes, RR=1.6 (95% CI 1.4 to 1.9), $p<0.001$).

Illnesses were reported from a variety of sports. In athletics, beach volleyball, football, sailing, synchronised swimming and taekwondo, just over 10% of the athletes suffered from at least one illness (table 1). The peak in relative illness rates occurred in beach volleyball, where 19% of the players suffered from an illness during the Games.

Table 2 Rates of overall injuries and injuries leading to time loss (>7 days of estimated absence) in female and male athletes in the Olympic sports

Olympic sport	Female athletes			Male athletes		
	n	All injuries	Time loss >7 days	n	All injuries	Time loss >7 days
Archery	64	1 (2.6)	0 (0.0)	64	1 (1.6)	0 (0.0)
Athletics	991	164 (16.5)	29 (2.9)	1088	190 (17.5)	28 (2.6)
Aquatics						
Diving	68	4 (5.9)	0 (0.0)	68	5 (7.4)	1 (1.5)
Swimming	450	26 (5.8)	1 (0.2)	481	24 (5.0)	1 (0.2)
Synchronised swimming	104	14 (13.5)	0 (0.0)	–	–	–
Water polo	104	9 (8.7)	0 (0.0)	156	25 (16.0)	0 (0.0)
Badminton	78	11 (14.1)	3 (3.8)	86	15 (17.4)	1 (1.2)
Basketball	143	14 (9.8)	1 (0.7)	144	18 (12.5)	6 (4.2)
Beach volleyball	48	5 (10.4)	0 (0.0)	48	7 (14.6)	0 (0.0)
Boxing	36	7 (19.4)	0 (0.0)	247	18 (7.3)	1 (0.4)
Canoe slalom	21	1 (4.8)	0 (0.0)	62	1 (1.6)	0 (0.0)
Canoe sprint	90	3 (3.3)	0 (0.0)	159	4 (2.5)	0 (0.0)
Cycling						
BMX	16	5 (31.3)	0 (0.0)	32	10 (31.3)	2 (6.3)
MTB	29	7 (24.1)	1 (3.4)	47	9 (19.1)	1 (2.1)
Road	66	7 (10.6)	0 (0.0)	144	12 (8.3)	2 (1.4)
Track	69	1 (1.2)	0 (0.0)	98	4 (4.1)	0 (0.0)
Equestrian	77	4 (5.2)	1 (1.3)	122	4 (3.3)	1 (0.8)
Fencing	124	7 (5.6)	0 (0.0)	122	16 (13.1)	2 (1.6)
Football	220	99 (45.0)	6 (2.7)	289	78 (27.0)	5 (1.7)
Gymnastics						
Artistic	97	6 (6.2)	2 (2.1)	98	9 (9.2)	2 (2.0)
Rhythmic	96	8 (8.3)	0 (0.0)	–	–	–
Trampoline	16	1 (6.3)	0 (0.0)	16	1 (6.3)	0 (0.0)
Handball	171	45 (26.3)	10 (5.8)	178	31 (17.4)	6 (3.4)
Hockey	192	33 (17.2)	4 (2.1)	196	34 (17.3)	6 (3.1)
Judo	153	19 (12.4)	3 (2.0)	230	25 (10.9)	8 (3.5)
Modern pentathlon	36	1 (2.8)	0 (0.0)	36	4 (11.1)	0 (0.0)
Rowing	196	8 (4.1)	0 (0.0)	353	10 (2.8)	0 (0.0)
Sailing	143	22 (15.4)	1 (0.7)	237	33 (13.9)	0 (0.0)
Shooting	159	11 (6.9)	0 (0.0)	231	4 (1.7)	0 (0.0)
Table tennis	88	3 (3.4)	0 (0.0)	86	8 (9.3)	2 (2.3)
Taekwondo	64	17 (26.6)	2 (3.1)	64	33 (51.6)	5 (7.8)
Tennis	89	9 (10.1)	2 (2.2)	95	12 (12.6)	2 (2.1)
Triathlon	55	10 (18.2)	1 (1.8)	55	6 (10.9)	2 (3.6)
Volleyball	144	11 (7.6)	3 (2.1)	144	9 (6.3)	0 (0.0)
Weightlifting	103	16 (15.5)	5 (4.9)	149	27 (18.1)	6 (4.0)
Wrestling	76	10 (13.2)	2 (2.6)	267	28 (10.5)	4 (1.5)
Total	4676	621 (13.3) *	76 (1.6)	5892	713 (12.1) *	94 (1.6)

Values are numbers (percentages) of injured athletes, unless otherwise indicated.

*Unknown gender for 27 injuries.

Affected system, causes, symptoms and severity of illness

A total of 310 illnesses (41%) affected the respiratory system, and these were most frequently observed in athletics, beach volleyball, football, swimming and water polo (3–5% of the athletes). In beach volleyball, 61% of the illnesses were reported to be respiratory infections. The second and third most affected systems were the gastrointestinal tract (n=123, 16%) and dermatology (n=83, 11%), respectively.

Concomitantly, infection was the most common cause of illness (n=347, 46%), affecting athletes in mainly the same sports as mentioned above. Symptoms of pain were present in 209 (16%) of the illnesses. However, for a third (248, 33%) of all illnesses, symptoms were not reported.

Almost one in five illnesses (n=145, 19%) were expected to result in absence from training or competition. Of these, nine illnesses were expected to result in an estimated time loss of more than 7 days (7 cases of stress, fatigue and exhaustion; 1 upper respiratory tract infection and 1 instance of abdominal pain).

Response rate and coverage of the athletes

Seventy-four of the 204 NOCs had more than 30 participating athletes. Athletes from these NOCs comprised 9450 of the total 10 568 athletes, corresponding to 89% (table 3). Throughout the 17 days of the London Games, the 74 NOCs submitted a

Table 3 Response rates, injuries and illnesses in NOCs of different sizes (measured by number of athletes)

	<30	30–99	100–200	>200	All
NOC (athletes)	132 (1114)	37 (1890)	20 (2510)	15 (5050)	204 (10 564*)
Injuries (%)	256 (23)	308 (16)	380 (15)	416 (8)	1360† (13)
Illnesses (%)	108 (10)	187 (10)	182 (7)	280 (6)	757‡ (7)
Report forms submitted§ (%)	–	624 (97)	332 (93)	248 (97)	1204 (96)
Injuries/illnesses reported by NOCs	43/18	163/105	238/117	291/190	735/430
Injuries/illnesses reported only by LOCOG	213/90	145/82	142/65	125/90	625/327

*Independent Olympic Athletes excluded.

†NOC is missing for 1 injury.

‡NOC is missing for 1 illness.

§Countries with less than 30 athletes were excluded from the response rate analysis.

LOCOG, London Organising Committee of the Olympic and Paralympic Games; NOCs, National Olympic Committee.

total of 1204 of a maximum of 1258 forms (mean 96%, range 12–100%).

A total of 625 injuries (46%) and 327 illnesses (43%) were missed by the NOCs and only recorded by the LOCOG medical staff. Of the 174 severe injuries, meaning those with an estimated absence for more than 7 days, almost three-quarters were captured by the NOC (n=123, 71%), while 51 injuries (29%) were reported solely through the medical encounters by the LOCOG medical staff.

DISCUSSION

This paper analyses and discusses the risk of injuries and illnesses among athletes competing in the London Olympic Games. The main findings of this 17-day long investigation were that 11% and 7% of all the 10 568 athletes suffered from at least one injury or illness, with overall rates of 128.8 injuries and 71.7 illnesses per 1000 athletes, respectively. The magnitude and characteristics of the injuries and illnesses varied substantially between sports and gender. While taekwondo, football, BMX and handball had the highest rates of injury, the highest rates of illness were found in beach volleyball, synchronised swimming and football.

Injury incidences in thelympic sports

The injury incidences found across the sports in the London Olympic Games are comparable to those reported earlier from the Athens Olympic Games in 2004 (team sports only),⁴ the Olympic Games in Beijing in 2008,¹⁸ as well as in certain major elite single-sport events.¹² However, the London Games athletes competing in archery or boxing experienced a 5% lower injury rate than their counterparts from the Beijing Olympic Games.¹⁸ Conversely, a higher injury rate was found among the London Games athletes competing in athletics (6%), badminton (11%), diving (6%), fencing (7%), handball (5%), sailing (14%), synchronised swimming (12%), taekwondo (12%), tennis (6%) and triathlon (5%). Similarly, the overall injury rate in athletics was 4–8% higher than that previously reported from the athletics World Championships from 2007 to 2012.^{13–15}

A change in injury incidence can be the result of changes in competition rules, in equipment or in other environmental factors. Changes in injury rates can also follow an increased or reduced awareness among both the athletes and their medical staff in recognising and reporting even minor incidents (broad injury and illness definition applied in IOC surveillance studies). In certain sports, changes may also be attributable to more comprehensive and accurate data reporting by team physicians who over time have been trained as injury and illness recorders

through the implementation of surveillance systems by their own Federation. Also, rate differences (lower or higher) may simply be the result of a natural fluctuation/variability of athletes exposure to risk, an observation that emphasises the value of on-going surveillance to monitor trends over time, for example, the effect of rule or equipment changes in the period between major sports events.

Severity, location and type of injuries

In the Olympic Games even minor injuries and illnesses with no or minimal time-loss can have serious consequences for the athlete. An injury or illness of otherwise insignificant severity may prevent Olympic athletes from reaching their life-time achievement. The majority (65%) of the injuries occurring in the London Olympic Games were reported to be of minimal severity, with no absence from competition or training. In contrast, 35% of all injuries were estimated to entail time-loss of at least 1 day from training or competition, while 13% were estimated to have more than 7 days time-loss. This indicates a reduction in the number of time-loss injuries from the Beijing Games, where half of the injuries were expected to prevent the athlete from participating in competition or training.¹⁸ Furthermore, of the more severe injuries leading to an absence of more than 7 days, 81% occurred during competition, supporting earlier findings that the injury rate is higher in competition than in training.^{8 13 15 21} In addition, it might indicate the overall importance of Olympic competition over training; the former for many athletes representing the culmination of their perennial efforts, sacrifices and preparation to fulfil their ambitions as Olympians.

The sports with the highest rate of injuries entailing a prolonged absence from training or competition (>7 days) were taekwondo (6% of the athletes), handball (5%), BMX cycling (4%) and weightlifting (4%). This corresponds with the data from the Beijing Games.

The risk of concussion is a recurrent concern in certain sports, and its diagnosis, prevention, treatment and return-to-play guidelines have been addressed in recent consensus meetings.^{22 23} However, as in the Beijing Games (12 concussions, 1.09 per 1000 registered athletes), the risk of sustaining a concussion in the London Games was low (6 concussions, 0.57 per 1000 athletes). In contrast, 20 concussions (7.8 per 1000 participating athlete) were reported from the 2010 Vancouver Winter Olympic Games, with every fifth injury overall affecting the head, neck and cervical spine.¹⁹ Pending more data, this may indicate that winter sports athletes, particularly those exposed to high speed are at even higher risk of potentially

severe head injuries than their summer-sports counterparts, including those competing in the martial arts.

Mechanisms and circumstances of injury

The mechanisms and circumstances of injuries in training and competition differed significantly among the different sports. Overall, the distribution of injuries across competition and training (55% vs 45%) was similar to that reported in the 2011 world championships in athletics,¹⁵ but more proportionate than those reported from the Beijing Games (74% vs 26%, respectively) and from other single-sport events.^{8–13} The majority of injuries were reported to be acute, whereas overuse injuries with either a gradual or sudden onset accounted for a quarter of the injuries. Although a similar distribution was reported from the Games in Beijing, these numbers should be interpreted with caution, owing to the current limitations in the recording of overuse injuries.²⁴

Illness risk during the Olympics

The impact of an illness on an athlete's competition and training can be just as significant as that of an injury. In recognition of this, the IOC and other major International Sports Federations, such as FIFA, FINA (Fédération Internationale de Natation) and IAAF (International Association of Athletics Federations) have extended their athlete health surveillance to also include illnesses.

Interestingly, the rate of illness in the London Games was identical to that of the Vancouver Winter Games, with 7 of 100 athletes being affected during the Games period. Similarly, sport-specific illness rates were comparable to those reported previously in aquatics (7% in London vs 7% in recent championships),¹² athletics (11% vs 7%)¹⁵ and football (12% vs 12%).⁸ Also consistent with the Vancouver data is the difference in the incidence of illnesses between female and male athletes, with women experiencing an illness incidence 60% higher than men. The same disproportion has previously been reported in the 2009 athletics¹³ and aquatics¹² world championships, but not in the 2011 athletics world championships¹⁵ or during the 1994–2009 US Open tennis championships.²⁵

The high incidence of respiratory infections mirrors data from other elite sport events.^{8–12–13–15–25} Predominant risk factors are mechanical and dehydration stresses generated within the airways and the level of airborne pollutants, irritants and allergens inhaled by the athlete under high ventilatory exercise conditions.²⁶ It has earlier been reported that airway hyper-responsiveness/asthma is the most common chronic medical condition experienced by both summer and winter Olympic athletes.²⁷

Methodological considerations

In studies on sports injury, it is usually recommended to express incidences using time exposed to risk as the denominator.^{20–28} However, given the inherent complexity of the Olympic Games, this was rendered not feasible in the present study. Instead we expressed the incidence of injury or illnesses as the number of new cases per 1000 athlete participants. Although this is the recommended approach in multisport events such as the Olympic Games,¹⁷ it ignores the fact that a multisport event is comprised of different frequencies and lengths of competition for each sport. For example, an Olympic diver's exposure to competition and training—and the encompassing risk of injury or illness—is by nature different to that of a sailor, weightlifter or footballer. Although interpretation of differences in injury incidence or pattern therefore should be made with some

caution, our reported incidences do indicate athletes' actual risk of suffering injuries or illnesses in their respective sport.

We defined injuries and illnesses as new or recurring injuries or illnesses receiving medical attention, regardless of the consequences with respect to absence from competition or training. By using such a definition, predominantly the moderate and severe acute injuries will be recorded. The less-serious injuries may be overlooked, since such injuries do not always require medical attention,^{29–30} albeit our results show that the majority of reported injuries were not estimated to entail any absence from the sport. In the Olympic Games, all athletes can get healthcare through the athletes' village polyclinic and the venue medical clinics. However, the availability, size and quality of the NOCs own medical teams vary between countries, meaning that not all athletes benefit from the same easy access to healthcare, which may bias the injury and illness recording. Nonetheless, we believe that our collection of data through both the NOCs and LOCOG physicians, we captured most injuries and illnesses.

Throughout the 17 days of data collection in the Olympic Games, we collected 96% of all the NOC injury and illness report forms. This is the highest NOC response rate to date in the Olympic Games injury and illness surveillance, a result which can be attributed to favourably disposed NOC medical staff, an informational meeting with all NOCs the day before the Games' opening, the preparation of report forms in eight languages and an instructional booklet on how to fill in the forms, and a dedicated research team which conducted frequent follow-up visits to boost NOCs' compliance. However, although the NOC response rate was very high, we did not test the accuracy and internal validity of their reported data. Thus, we cannot know the extent to which the NOC data match the actual circumstances of the occurred injury or illness. An indication of this is that a number of injuries and illnesses were recorded in an incomplete manner with missing information in categories such as type, location and diagnosis. Furthermore, it has been documented previously in professional alpine skiing¹⁶ and male elite football³¹ that prospective injury surveillance by team medical staff underestimates the incidence of injuries and time-loss injuries. Our results support these findings; compared with the medical staff working at the polyclinic and medical venues, the NOC medical teams did not report 46% of the injuries, 29% of the time-loss injuries and 43% of the illnesses. With respect to the more severe injuries the results were somewhat better; of the 174 injuries with an estimated absence of more than 7 days, almost three-quarters were captured by the NOCs. Retrospective athlete interviews at the end of the event have been suggested as one option to improve the surveillance data quality¹⁵; however, this seems impractical in major events comprising thousands of athletes, such as the Olympic Games.

Recent papers from Bahr²⁴ and Clarsen *et al*^{32–33} highlight the inadequacy of the current standard epidemiological methodology to record the true magnitude and pattern of overuse injuries in sport. As pointed out by Bahr,²⁴ overuse injuries in many sports represent as much of a problem as do acute injuries, but these injuries are seldom properly captured or registered in statistics through current recording methods, as athletes with overuse problems often do not seek medical attention or lose time from training or competition. In the present study, overuse was reported to be the mechanism for every fourth injury, notably in diving, athletics, weightlifting, swimming and fencing. Furthermore, 68% of the overuse injuries in London were recorded with no estimated absence, an indication that—also in the Olympic setting—athletes with overuse problems often continue to train and compete, although typically with

significant inhibitions, such as pain, restricted function and reduced performance.

Practical implications

The continuous surveillance of injury and illness rates is a fundamental pre-requisite to athlete health protection. Only by monitoring the development over several years can we identify and follow changes in potential risk factors and mechanisms of injury and illnesses in the different sports and their disciplines, which, in turn, will allow us to develop, introduce and update targeted preventive measures. The key to a successful study of epidemiology lies in a well-organised data-collection procedure with coordinated efforts from sports medicine professionals, coaches and athletes, combined with systematic subsequent analyses.

However, surveillance of injuries and illnesses for elite athletes during multisport events such as the Olympic Games is subject to other challenges than surveillance systems in other contexts, such as single sport or amateur events. Differences encompass: the limited data-collection window; the great amount of data to be recorded and processed very quickly; the vast array of people—with different professional backgrounds—involved in the same data-recording process; the high number of different specific settings in which the injuries occur (eg, canoeing vs tennis vs taekwondo); the low level of severe health incidents, with most being minor-to-moderate musculoskeletal conditions and the intense media and public scrutiny of injuries and illnesses when the athletes fail to meet their performance goals.³⁴ The IOC injury and illness surveillance system has been continuously improved to best tackle these unique conditions, and we aim to continue to further refine its processes.

For example, a long-term goal of the IOC is to develop a customised electronic health record (EHR) that (1) can be used by NOCs on both a general year round basis in the clinic to record and follow athletes' health/injury, employing periodic health evaluations (PHE) recommendations,³⁵ and in the lead up to and during the Olympic Games; (2) can be used at the Olympic Games' polyclinics and venue medical centres to have an immediate medical history of the athlete, which can be taken into consideration to optimise the treatment of injuries and illnesses during the Olympic Games and (3) will provide the IOC and other users with longitudinal de-identified individual and aggregate data allowing the IOC to survey and assess the epidemiology, identify injury and illness risk factors and mechanisms, and in turn, introduce tailored measures to prevent injuries and illnesses in athletes in all sports. We believe that instituting an EHR in the Olympic Movement can have a tremendous impact on athletes' health protection. Furthermore, it would also remedy the current recording problem of overuse injuries. By adopting the recommendations of Bahr²⁴ and Clarsen *et al*^{32 33} the EHR would facilitate the prospective serial measurement and recording of athlete health and symptoms of pathology, using sensitive scoring instruments for pain and other overuse symptoms, and open up for reporting of both the incidence and prevalence of injuries.

The continuously accumulating evidence that injury and illness rates vary substantially between sports demonstrates the need for tailoring preventive measures to the specific context of each sport. Sport bodies such as the IOC, IFs and NOCs have a great responsibility to protect the health of their athletes. The Olympic Movement Medical Code encourages all stakeholders to take measures to ensure that sport is practised without danger to the health of the athletes and to minimise the risks of physical injury and psychological harm. For NOCs one evident

way of achieving this is by introducing PHEs in their daily practice, which are instrumental in preventing injuries and illnesses, and hence, to protect the health of the athletes. For IFs, a critical component of this responsibility is the institution of a scientifically sound injury and illness surveillance system in all major events. Some sports federations, such as FIFA, FINA, FIS (Fédération Internationale de Ski), FIVB (Fédération Internationale de Volleyball), IAAF and UEFA (Union of European Football Associations) have put increasing efforts into working systematically and scientifically to protect their athletes' health. We encourage other IFs and sports organisations to follow their example.

CONCLUSION

Eleven per cent of the athletes incurred an injury during the London Olympic Games, and 7% suffered from at least one illness. The incidences and characteristics of injuries and illnesses during training and competition varied substantially between sports and gender. Future initiatives should include the development of preventive measures tailored for each specific sport and the continued focus among sport bodies to institute and further develop scientific injury and illness surveillance systems.

Author affiliations

¹Medical & Scientific Department, International Olympic Committee, Lausanne, Switzerland

²Department of Orthopaedic Surgery, University of Oslo, Oslo, Norway

³Department of Sports Medicine, Oslo Sports Trauma Research Center, Norwegian School of Sport Sciences, Oslo, Norway

⁴International Association of Athletics Federations (IAAF), Monte Carlo, Monaco

⁵International Ice Hockey Federation (IIHF), Zurich, Switzerland

⁶Fédération Internationale de Football de Football Association (FIFA), Zurich, Switzerland

⁷FIFA Medical Assessment and Research Center (F-MARC) and Schulthess Klinik, Zurich, Switzerland

⁸Olympic Council of Asia (OCA), Hawalli, Kuwait

⁹Commonwealth Games Federation (CGF), London, UK

¹⁰Olympic Council of Malaysia (MAS), Kuala Lumpur, Malaysia

¹¹Sport Injury Prevention Research Centre, University of Calgary, Alberta, Canada

¹²Fédération Internationale de Natation (FINA), Lausanne, Switzerland

¹³Division of Orthopaedic and Accident Surgery, University of Nottingham, Nottingham, UK

¹⁴University College London, London, UK

¹⁵Stockholm Sports Trauma Research Center, Karolinska Institutet, Stockholm, Sweden

Acknowledgements The authors specially thank Professor Fares Haddad for his assistance with obtaining the UCL Research Ethics Committee approval. The authors also sincerely thank all the NOC medical staff contributing to the data collection: Dr Bensoultane Zaher (ALG), Dr Maria Stella Cristiano (ANG), Dr Hugo Rodriguez (ARG), Dr Peter Baquie and colleagues (AUS), Dr Alfred Engel (AUT), Dr Adrian Lorde (BAR), Dr Marc Vangrinsven (BEL), Dr Yauheni Lasitski (BLR), Dr Jose Alfredo Padilha (BRA), Dr Stefan Strugarov (BUL), Dr Robert McCormack (CAN), Dr Guoping Li (CHN), Dr Boubakary Sidiki (CMR), Dr Mauricio Serrato Rua (COL), Dr Boris Labar (CRO), Dr Jorge Pavel Pinos Rivero (CUB), Dr Vetvicka and Dr Heinz (CZE), Dr Morten Storgaard (DEN), Dr Milton Pineda (DOM), Dr Flores and Dr Eimen (ECU), Dr Mostafa Elmofly (EGY), Dr Ramon Olive (ESP), Dr Mihkel Mardna (EST), Dr Ayalew Tilahun Beshaha (ETH), Dr Tommi Vasankari (FIN), Dr Philippe Levan (FRA), Dr Eric Baye (GAB), Dr Ian McCurdie and colleagues (GBR), Zurab Kakhabrishvili (GEO), Dr Bernd Wolfarth (GER), Dr Angela El-Adas (GHA), Dr Georgios Marinou and colleagues (GRE), Dr Hans Larsen (HAI), Dr Julian Wai Chang (HKG), Dr Eva Villalta Rivera (HON), Dr Peter Balogh (HUN), Dr Sarla Rao (IND), Dr Lotfali Pourkazemi (IRI), Dr Sean Gaine and colleagues (IRL), Dr Iftach Hetsroni (ISR), Dr Antonio Pelliccia (ITA), Dr Warren Blake (JAM), Dr Takao Akama and colleagues (JPN), Dr Sagit Bavaev (KAZ), Dr Julius Ogeto (KEN), Dr Won Hah Park and colleagues (KOR), Dr Abdullah Ashour (KSA), Dr Liga Cirule (LAT), Dr German Clénin (LIE), Dr Dalius Barkauskas (LTU), Dr Axel Urhausen (LUX), Dr Abdellali Sobhi (MAR), Dr Ramlan Bin Abd Aziz (MAS), Dr Hugo Martinez (MEX), Dr Lucienne Attard (MLT), Dr Predag Dabovic (MNE), Dr Barend Phillipus 'Ben' Viljoen (NAM), Dr Abd-Rein van den Hoogenband (NED), Dr Abdulkadir Mu'azu and Dr Ibrahim Gyaran (NGR), Dr Tonje Wåle Flørenes (NOR), Dr Lynne Coleman and colleagues

(NZL), Dr Caleb Otto (PLW), Dr Hubert Krysztofiak and colleagues (POL), Dr Artur Teixeira Pergira De Castro (POR), Dr Kim Yong Thaek and colleagues (PRK), Dr Jose Correa (PUR), Dr Rachid Bouras (QAT), Dr Dan Florin Tanase (ROU), Dr Phatho Cele and colleagues (RSA), Dr Maxim Velichko (RUS), Dr Abba Diatta (SEN), Dr Ahtime and Dr Sherwin (SEY), Dr Cormac O'Muircheartaigh (SIN), Dr Dragan Radovanic (SRB), Dr German Clénin (SUI), Dr Branislav Delej (SVK), Dr Bo Berglund (SWE), Dr Hilary Meechai Inwood (THA), Dr Chou and Dr Chang (TPE), Dr Terry ALI (TRI), Dr Haj Romdhane M. R. and Dr Moncef Ben Abid (TUN), Dr Tugba Kocahan (TUR), Dr Abdulhameed Al Attar (UAE), Dr Olesandr Varvinskyi (UKR), Dr Gonzalo Andrés Rodríguez Gómez (URU), Dr Chang and Dr Moreau (USA), Dr Elena Abaeva (UZB), Dr Marelia Guillen Mujica (VEN), Dr Nguyen Van Phu (VIE), Dr Nicholas Munyonga (ZIM).

Competing interests None.

Provenance and peer review Commissioned; internally peer reviewed.

REFERENCES

- International Olympic Committee. Olympic Charter—the organisation, action and operation of the Olympic Movement. 2011. http://www.olympic.org/Documents/olympic_charter_en.pdf (accessed Feb 2013).
- Van Mechelen W, Hlobil H, Kemper HC. Incidence, severity, aetiology and prevention of sports injuries. A review of concepts. *Sports Med* 1992;14:82–99.
- Junge A, Dvorak J, Graf-Baumann T, et al. Football injuries during FIFA tournaments and the Olympic Games, 1998–2001: development and implementation of an injury-reporting system. *Am J Sports Med* 2004;32(Suppl 1):805–95.
- Junge A, Langevoort G, Pipe A, et al. Injuries in team sport tournaments during the 2004 Olympic Games. *Am J Sports Med* 2006;34:565–76.
- Junge A, Dvorak J, Graf-Baumann T. Football injuries during the World Cup 2002. *Am J Sports Med* 2004;32(Suppl 1):235–75.
- Yoon YS, Chai M, Shin DW. Football injuries at Asian tournaments. *Am J Sports Med* 2004;32(Suppl 1):365–425.
- Junge A, Dvorak J. Injuries in female football players in top-level international tournaments. *Br J Sports Med* 2007;41(Suppl 1):i3–7.
- Dvorak J, Junge A, Derman W, et al. Injuries and illnesses of football players during the 2010 FIFA World Cup. *Br J Sports Med* 2011;45:626–30.
- Bahr R, Reeser JC. Injuries among world-class professional beach volleyball players. The Federation Internationale de Volleyball beach volleyball injury study. *Am J Sports Med* 2003;31:119–25.
- Langevoort G, Myklebust G, Dvorak J, et al. Handball injuries during major international tournaments. *Scand J Med Sci Sports* 2007;17:400–7.
- Fuller CW, Laborde F, Leather RJ, et al. International Rugby Board Rugby World Cup 2007 injury surveillance study. *Br J Sports Med* 2008;42:452–9.
- Mountjoy M, Junge A, Alonso JM, et al. Sports injuries and illnesses in the 2009 FINA World Championships (Aquatics). *Br J Sports Med* 2010;44:522–7.
- Alonso JM, Tscholl PM, Engebretsen L, et al. Occurrence of injuries and illnesses during the 2009 IAAF World Athletics Championships. *Br J Sports Med* 2010;44:1100–5.
- Alonso JM, Junge A, Renström P, et al. Sports injuries surveillance during the 2007 IAAF World Athletics Championships. *Clin J Sport Med* 2009;19:26–32.
- Alonso JM, Edouard P, Fischetto G, et al. Determination of future prevention strategies in elite track and field: analysis of Daegu 2011 IAAF Championships injuries and illnesses surveillance. *Br J Sports Med* 2012;46:505–14.
- Flørenes TW, Nordsletten L, Heir S, et al. Recording injuries among World Cup skiers and snowboarders: a methodological study. *Scand J Med Sci Sports* 2011;21:196–205.
- Junge A, Engebretsen L, Alonso JM, et al. Injury surveillance in multi-sport events: the International Olympic Committee approach. *Br J Sports Med* 2008;42:413–21.
- Junge A, Engebretsen L, Mountjoy ML, et al. Sports injuries during the Summer Olympic Games 2008. *Am J Sports Med* 2009;37:2165–72.
- Engebretsen L, Steffen K, Alonso JM, et al. Sports injuries and illnesses during the Winter Olympic Games 2010. *Br J Sports Med* 2010;44:772–80.
- Fuller CW, Ekstrand J, Junge A, et al. Consensus statement on injury definitions and data collection procedures in studies of football (soccer) injuries. *Br J Sports Med* 2006;40:193–201.
- Møller M, Attermann J, Myklebust G, et al. Injury risk in Danish youth and senior elite handball using a new SMS text messages approach. *Br J Sports Med* 2012;46:531–7.
- McCroory P, Meeuwisse W, Johnston K, et al. Consensus statement on concussion in sport: the 3rd international conference on concussion in sport held in Zurich, November 2008. *Br J Sports Med* 2009;43(Suppl 1):i76–90.
- McCroory P, Meeuwisse W, Aubry M, et al. Consensus statement on concussion in sport—the 4th international conference on concussion in sport held in Zurich, November 2012. *Br J Sports Med* 2013;47:250–8.
- Bahr R. No injuries, but plenty of pain? On the methodology for recording overuse symptoms in sports. *Br J Sports Med* 2009;43:966–72.
- Sell K, Hainline B, Yorio M, et al. Illness data from the US Open tennis championships from 1994 to 2009. *Clin J Sport Med* 2013;23:25–32.
- Kippelen P, Fitch KD, Anderson SD, et al. Respiratory health of elite athletes—preventing airway injury: a critical review. *Br J Sports Med* 2012;46:471–6.
- Fitch K, Anderson S. Intense exercise and airway hyper-responsiveness/asthma: importance of environmental factors. *Br J Sports Med* 2012;46:379–80.
- Bahr R, Holme I. Risk factors for sports injuries—a methodological approach. *Br J Sports Med* 2003;37:384–92.
- Inklaar H. Soccer injuries. I: incidence and severity. *Sports Med* 1994;18:55–73.
- Finch CF. An overview of some definitional issues for sports injury surveillance. *Sports Med* 1997;24:157–63.
- Bjørneboe J, Flørenes TW, Bahr R, et al. Injury surveillance in male professional football; is medical staff reporting complete and accurate? *Scand J Med Sci Sports* 2010;21:713–20.
- Clarsen B, Myklebust G, Bahr R. Development and validation of a new method for the registration of overuse injuries in sports injury epidemiology: the Oslo Sports Trauma Research Centre (OSTRC) overuse injury questionnaire. *Br J Sports Med* 2013. Published online first: 8 February 2013. <http://www.ncbi.nlm.nih.gov/pubmed/23038786>
- Clarsen B, Rønsen O, Myklebust G, et al. The Oslo Sports Trauma Research Center questionnaire on health problems: a new approach to prospective monitoring of illness and injury in elite athletes. *Br J Sports Med* 2013. Published online first: 21 February 2013. <http://www.ncbi.nlm.nih.gov/pubmed/23429267>.
- Finch CF. Injury surveillance, the Olympic way. 2012. <http://blogs.bmj.com/injury-prevention/2012/07/30/injury-surveillance-the-olympic-way/> (accessed Feb 2013).
- Ljungqvist A, Jenoure P, Engebretsen L, et al. The International Olympic Committee (IOC) consensus statement on periodic health evaluation of elite athletes March 2009. *Br J Sports Med* 2009;43:631–43.